



LEQEMBI IQLIK® (lecanemab-irmb) Autoinjector for Initiation of Therapy Now Available in the U.S. for Early Alzheimer's Disease

August 24, 2026

LEQEMBI IQLIK is the first-of-its-kind anti-amyloid treatment worldwide offering at-home dosing for initiation and maintenance (approved in the U.S.)

TOKYO and CAMBRIDGE, Mass., Aug. 24, 2026 (GLOBE NEWSWIRE) -- Eisai Co., Ltd. and Biogen Inc. (Nasdaq: BIIB), announced today that once weekly lecanemab-irmb subcutaneous injection (brand name: LEQEMBI IQLIK®) is now available in the U.S. for initiation therapy for early Alzheimer's disease (AD) in adults with mild cognitive impairment (MCI) or mild dementia due to AD, collectively referred to as early AD. MCI due to AD is the earliest symptomatic stage of AD and can appear with subtle symptoms such as forgetfulness, confusion, or feeling at a loss for words.¹

LEQEMBI IQLIK is administered via an autoinjector, introducing a convenient alternative to intravenous (IV) dosing from the start of treatment. For initiation, the approved regimen is 500 mg given once weekly as two consecutive 250 mg injections, each delivered in approximately 15 seconds. LEQEMBI IQLIK may also be used for maintenance dosing at 360 mg once weekly after 18 months of IV or subcutaneous (SC) treatment. Throughout the entire treatment course – from initiation through maintenance – patients may receive LEQEMBI either as IV infusion or as SC injection with LEQEMBI IQLIK. Patients may also switch from IV to SC administration, or vice versa, providing greater convenience and flexibility in LEQEMBI administration.

Expanding Treatment Convenience and Flexibility Across the Alzheimer's Disease Care Pathway

The availability of LEQEMBI IQLIK for both initiation and maintenance therapy in the U.S. enhances treatment convenience and flexibility, offering early AD patients and their care partners more control in managing their care while lowering barriers to initiating and continuing treatment with LEQEMBI. LEQEMBI IQLIK may reduce the time spent receiving anti-amyloid therapy via IV infusions. In addition, at-home administration allows patients and their care partners to continue treatment without the burden of clinic visits, making it easier to go out and travel. LEQEMBI IQLIK also has the potential to reduce healthcare resources associated with IV dosing, such as infusion preparation and nurse monitoring. These features may help streamline the overall AD treatment pathway. For ARIA monitoring, as with IV administration, brain magnetic resonance imaging (MRI) is performed prior to initiating treatment and at specified time points after treatment initiation.

Support Resources for LEQEMBI IQLIK

Eisai and Biogen will provide a range of resources to help patients, care partners, healthcare providers, health systems and pharmacies successfully implement and use LEQEMBI IQLIK.

Patient Support

Resources include an Instructions for Use (IFU) video and IQLIK Welcome Kit. The IQLIK Welcome Kit provides a What to Expect Treatment Tracker, an injection reminder magnet, and a Demo Kit to help patients and care partners understand what to expect, prepare for at-home injections, and administer injections safely at home. The LEQEMBI Companion® app will also be available. It brings the information patients and care partners need into one experience, including education about the injection process and tools for tracking each dose.

Patients initiating IQLIK will be encouraged to bring the Welcome Kit to their initial dosing appointment. Kits will be available through:

- The LEQEMBI Specialty Pharmacy Network (CareMed Specialty Pharmacy and Walgreens Specialty Pharmacy)
- Integrated Delivery Networks (IDNs) or the prescribing physician
- LEQEMBI Companion® - a program offering financial support information and options for eligible patients.

Through the LEQEMBI Specialty Pharmacy Network, patients may receive support with prescription fulfillment, insurance coverage navigation, onboarding, delivery coordination, device-use education, and – if they choose to participate – treatment reminders and educational support during the first six months of therapy. Patients using other eligible specialty pharmacies may have access to similar support, which may vary by pharmacy.

Healthcare Provider Resources

Electronic Health Record (EHR) support materials and a step-by-step Getting Started Guide will help providers navigate the process from prescription submission through therapy initiation.

Financial Assistance

To further support access to LEQEMBI for certain patients who need help paying for their medicines, Eisai's Patient Assistance Program (PAP) will provide LEQEMBI and LEQEMBI IQLIK at no cost, for eligible uninsured patients, who meet financial need and other program criteria.

*The LEQEMBI Companion app was developed to deliver behavior-driven digital support for patients and care partners along their treatment journey, in partnership with Medisafe Ltd., a behavioral patient engagement engine built for complex and specialty therapies. Patients can visit LEQEMBI.com/CompanionAppSignUp to get started.

Eisai serves as the lead for lecanemab's development and regulatory submissions globally with Eisai and Biogen co-commercializing and co-promoting the product and Eisai having final decision-making authority.

INDICATION

LEQEMBI® is indicated for the treatment of Alzheimer's disease (AD). Treatment with LEQEMBI should be initiated in patients with mild cognitive impairment (MCI) or mild dementia stage of disease, the population in which treatment was initiated in clinical trials.

IMPORTANT SAFETY INFORMATION

WARNING: AMYLOID-RELATED IMAGING ABNORMALITIES (ARIA)
--

- Monoclonal antibodies directed against aggregated forms of beta amyloid, including LEQEMBI, can cause ARIA, characterized as ARIA with edema (ARIA-E) and ARIA with hemosiderin deposition (ARIA-H). Incidence and timing of ARIA vary among treatments. ARIA usually occurs early in treatment and is usually asymptomatic, although serious and life-threatening events, including seizure and status epilepticus, can occur. ARIA can be fatal. Serious intracerebral hemorrhages (ICH) >1 cm, some of which have been fatal, have been observed with this class of medications. Because ARIA-E can cause focal neurologic deficits that can mimic an ischemic stroke, consider whether such symptoms could be due to ARIA-E before giving thrombolytic therapy to a patient being treated with LEQEMBI.
 - **Apolipoprotein E ε4 (ApoE ε4) Homozygotes:** Patients who are ApoE ε4 homozygotes (~15% of patients with AD) treated with this class of medications have a higher incidence of ARIA, including symptomatic, serious, and severe radiographic ARIA, compared to heterozygotes and noncarriers. Testing for ApoE ε4 status should be performed prior to initiation of treatment to inform the risk of developing ARIA. Prior to testing, prescribers should discuss with patients the risk of ARIA across genotypes and the implications of genetic testing results. Prescribers should inform patients that if genotype testing is not performed, they can still be treated with LEQEMBI; however, it cannot be determined if they are ApoE ε4 homozygotes and at higher risk for ARIA.
- Consider the benefit of LEQEMBI for the treatment of AD and the potential risk of serious ARIA events when deciding to initiate treatment with LEQEMBI.

CONTRAINDICATION

Contraindicated in patients with serious hypersensitivity to lecanemab-irmb or to any of the excipients. Reactions have included angioedema and anaphylaxis.

WARNINGS AND PRECAUTIONS

AMYLOID-RELATED IMAGING ABNORMALITIES

Medications in this class, including LEQEMBI, can cause ARIA-E, which can be observed on MRI as brain edema or sulcal effusions, and ARIA-H, which includes microhemorrhage and superficial siderosis. ARIA can occur spontaneously in patients with AD, particularly in patients with MRI findings suggestive of cerebral amyloid angiopathy (CAA), such as pretreatment microhemorrhage or superficial siderosis. ARIA-H generally occurs with ARIA-E. Reported ARIA symptoms may include headache, confusion, visual changes, dizziness, nausea, and gait difficulty. Focal neurologic deficits may also occur. Symptoms usually resolve over time.

Incidence of ARIA

Symptomatic ARIA occurred in 3% and serious ARIA symptoms in 0.7% with LEQEMBI. Clinical ARIA symptoms resolved in 79% of patients during the period of observation. ARIA, including asymptomatic radiographic events, was observed: LEQEMBI, 21%; placebo, 9%. ARIA-E was observed: LEQEMBI, 13%; placebo, 2%. ARIA-H was observed: LEQEMBI, 17%; placebo, 9%. No increase in isolated ARIA-H was observed for LEQEMBI vs placebo.

Incidence of ICH

ICH >1 cm in diameter was reported in 0.7% with LEQEMBI vs 0.1% with placebo. Fatal events of ICH in patients taking LEQEMBI have been observed.

Risk Factors of ARIA and ICH

ApoE ε4 Carrier Status

Of the patients taking LEQEMBI, 16% were ApoE ε4 homozygotes, 53% were heterozygotes, and 31% were noncarriers. With LEQEMBI, ARIA was higher in ApoE ε4 homozygotes (LEQEMBI: 45%; placebo: 22%) than in heterozygotes (LEQEMBI: 19%; placebo: 9%) and noncarriers (LEQEMBI: 13%; placebo: 4%). Symptomatic ARIA-E occurred in 9% of ApoE ε4 homozygotes vs 2% of heterozygotes and 1% of noncarriers. Serious ARIA events occurred in 3% of ApoE ε4 homozygotes and in ~1% of heterozygotes and noncarriers. The recommendations on management of ARIA do not differ between ApoE ε4 carriers and noncarriers.

Radiographic Findings of CAA

Neuroimaging findings that may indicate CAA include evidence of prior ICH, cerebral microhemorrhage, and cortical superficial siderosis. CAA has an increased risk for ICH. The presence of an ApoE ε4 allele is also associated with CAA.

The baseline presence of at least 2 microhemorrhages or the presence of at least 1 area of superficial siderosis on MRI, which may be suggestive of CAA, have been identified as risk factors for ARIA. Patients were excluded from Clarity AD for the presence of >4 microhemorrhages and additional findings suggestive of CAA (prior cerebral hemorrhage >1 cm in greatest diameter, superficial siderosis, vasogenic edema) or other lesions (aneurysm, vascular malformation) that could potentially increase the risk of ICH.

Concomitant Antithrombotic or Thrombolytic Medication

In Clarity AD, baseline use of antithrombotic medication (aspirin, other antiplatelets, or anticoagulants) was allowed if the patient was on a stable dose. Most exposures were to aspirin. Antithrombotic medications did not increase the risk of ARIA with LEQEMBI. The incidence of ICH: 0.9% in patients taking LEQEMBI with a concomitant antithrombotic medication vs 0.6% with no antithrombotic and 2.5% in patients taking LEQEMBI with an anticoagulant alone or with antiplatelet medication such as aspirin vs none in patients receiving placebo.

Fatal cerebral hemorrhage has occurred in 1 patient taking an anti-amyloid monoclonal antibody in the setting of focal neurologic symptoms of ARIA and the use of a thrombolytic agent.

Additional caution should be exercised when considering the administration of antithrombotics or a thrombolytic agent (e.g., tissue plasminogen activator) to a patient already being treated with LEQEMBI. Because ARIA-E can cause focal neurologic deficits that can mimic an ischemic stroke, treating clinicians should consider whether such symptoms could be due to ARIA-E before giving thrombolytic therapy in a patient being treated with LEQEMBI.

Caution should be exercised when considering the use of LEQEMBI in patients with factors that indicate an increased risk for ICH and, in particular, patients who need to be on anticoagulant therapy or patients with findings on MRI that are suggestive of CAA.

Radiographic Severity With LEQEMBI

Most ARIA-E radiographic events occurred within the first 7 doses, although ARIA can occur at any time, and patients can have >1 episode. Maximum radiographic severity of ARIA-E with LEQEMBI was mild in 4%, moderate in 7%, and severe in 1% of patients. Resolution on MRI occurred in 52% of

ARIA-E patients by 12 weeks, 81% by 17 weeks, and 100% overall after detection. Maximum radiographic severity of ARIA-H microhemorrhage with LEQEMBI was mild in 9%, moderate in 2%, and severe in 3% of patients; superficial siderosis was mild in 4%, moderate in 1%, and severe in 0.4% of patients. With LEQEMBI, the rate of severe radiographic ARIA-E was highest in ApoE ε4 homozygotes (5%) vs heterozygotes (0.4%) or noncarriers (0%). With LEQEMBI, the rate of severe radiographic ARIA-H was highest in ApoE ε4 homozygotes (13.5%) vs heterozygotes (2.1%) or noncarriers (1.1%).

Monitoring and Dose Management Guidelines

Baseline brain MRI and periodic monitoring with MRI are recommended. Enhanced clinical vigilance for ARIA is recommended during the first 14 weeks of treatment. Depending on ARIA-E and ARIA-H clinical symptoms and radiographic severity, use clinical judgment when considering whether to continue dosing or to temporarily or permanently discontinue LEQEMBI. If a patient experiences ARIA symptoms, clinical evaluation should be performed, including MRI if indicated. If ARIA is observed on MRI, careful clinical evaluation should be performed prior to continuing treatment.

HYPERSENSITIVITY REACTIONS

Hypersensitivity reactions, including angioedema, bronchospasm, and anaphylaxis, have occurred with LEQEMBI. Promptly discontinue the infusion upon the first observation of any signs or symptoms consistent with a hypersensitivity reaction and initiate appropriate therapy.

INFUSION-RELATED REACTIONS (IRRs)

IRRs were observed—LEQEMBI: 26%; placebo: 7%—and most cases with LEQEMBI (75%) occurred with the first infusion. IRRs were mostly mild (69%) or moderate (28%). Symptoms included fever and flu-like symptoms (chills, generalized aches, feeling shaky, and joint pain), nausea, vomiting, hypotension, hypertension, and oxygen desaturation.

IRRs can occur during or after the completion of infusion. In the event of an IRR during the infusion, the infusion rate may be reduced or discontinued, and appropriate therapy initiated as clinically indicated. Consider prophylactic treatment prior to future infusions with antihistamines, acetaminophen, nonsteroidal anti-inflammatory drugs, or corticosteroids.

ADVERSE REACTIONS

- The most common adverse reactions reported in ≥5% with LEQEMBI infusion every 2 weeks and ≥2% higher than placebo were IRRs (LEQEMBI: 26%; placebo: 7%), ARIA-H (LEQEMBI: 14%; placebo: 8%), ARIA-E (LEQEMBI: 13%; placebo: 2%), headache (LEQEMBI: 11%; placebo: 8%), superficial siderosis of central nervous system (LEQEMBI: 6%; placebo: 3%), rash (LEQEMBI: 6%; placebo: 4%), and nausea/vomiting (LEQEMBI: 6%; placebo: 4%)
- The safety profile of subcutaneous LEQEMBI was similar to intravenous infusion. Subcutaneous dosing was associated with mostly localized (erythema, induration, swelling, heat, pain, pruritus, rash, ecchymosis, nodule, and hematoma) and less frequent systemic (headache, chills, fever, and fatigue) injection-related reactions, majority at first dose when initiating therapy. Localized reactions that were recurrent and/or delayed were observed. Severe localized reactions and cases leading to dose discontinuation or interruption occurred.

LEQEMBI (lecanemab-irmb) is available:

- Intravenous infusion: 100 mg/mL
- Subcutaneous injection: 200 mg/mL

Please see full [Prescribing Information](#) for LEQEMBI, including **Boxed WARNING**.

Click [here](#) to access the LEQEMBI digital library with assets available for download.

MEDIA CONTACTS

Eisai Co., Ltd.

Public Relations Department
TEL: +81 (0)3-3817-5120

Eisai Europe, Ltd.

EMEA Communications Department
+44 (0) 7760 619251
Emea-comms@eisai.net

Eisai Inc. (U.S.)

Julie Edelman
+1-862-213-5915
Julie_Edelman@eisai.com

Biogen Inc.

Madeleine Shin
+1-781-464-3260
public.affairs@biogen.com

INVESTOR CONTACTS

Eisai Co., Ltd.

Investor Relations Department
TEL: +81 (0) 3-3817-5122

Biogen Inc.

Tim Power
+ 1-781-464-2442
IR@biogen.com

Notes to Editors

1. About lecanemab (generic name, brand name: LEQEMBI®)

Lecanemab is the result of a strategic research alliance between Eisai and BioArctic. It is a humanized immunoglobulin gamma (IgG1) monoclonal antibody directed against aggregated soluble (protofibril) and insoluble forms of amyloid-beta (Aβ).

Lecanemab has been approved in 53 countries and regions including Japan, the U.S., China, Europe, South Korea, Taiwan, and Saudi Arabia, and is under regulatory review in 6 countries. Following the initial phase with treatment every two weeks for 18 months, intravenous (IV) maintenance dosing

with treatment every four weeks is approved in 8 countries including the U.S., China, the UK, and others, and applications have been filed in 12 countries and regions. The U.S. FDA approved Eisai's Biologics License Application (BLA) for subcutaneous maintenance dosing with LEQEMBI IQLIK in August 2025. For subcutaneous initiation treatment (500 mg), approval was obtained in the United States in July 2026, and applications are under review in four countries, including Japan and China. In China, the application has been granted Priority Review designation. Since December 2025, lecanemab (IV) has been included in the "Commercial Insurance Innovative Drug List," recently introduced by the National Healthcare Security Administration (NHSA) of China.

LEQEMBI's approvals in these countries were based on Phase 3 data from Eisai's global placebo-controlled, double-blind, parallel-group, randomized Clarity AD clinical trial, in which it met its primary endpoint and all key secondary endpoints with statistically significant results. The primary endpoint was the global cognitive and functional scale, Clinical Dementia Rating Sum of Boxes (CDR-SB). Clarity AD evaluated lecanemab 10 mg/kg bi-weekly IV treatment of early Alzheimer's disease, which involved 1,795 patients (treatment group: 898, placebo group: 897). 95% of patients who completed the core study (18 months) chose to continue in the long-term extension study (LTE), with 478 patients still receiving treatment for four years. In the Clarity AD core clinical study, data showed LEQEMBI IV significantly slowed disease progression at 18 months (27% vs placebo), and the mean change from baseline between the lecanemab treated group and the placebo group after 18 months was -0.45 ($P=0.00005$) on the primary endpoint of CDR-SB global cognitive and functional scale.

To provide context, a change from 0.5 to 1 on the Clinical Dementia Rating (CDR) score domains of Memory, Community Affairs and Home/Hobbies reflects a shift from mild impairment to loss of independence. This can affect a person's ability to be left alone safely, recall recent events, participate in daily activities, manage household tasks, and engage in hobbies and intellectual interests.

LEQEMBI also rapidly reduced plaque as early as three months (-59.1 CL difference vs placebo in amyloid level at 18 months; $P<0.00001$).^{*} Additionally, LEQEMBI continued to show benefit over a four-year LTE treatment period; in a subgroup analysis, 81 percent of LEQEMBI patients who stayed on treatment remained in the early AD stages at four years.^{**}

Over three years of treatment, including both the core study and the LTE, data showed lecanemab demonstrated a reduction in cognitive decline—measured by CDR-SB—of 1.01 points compared to the expected decline observed in the Alzheimer's Disease Neuroimaging Initiative (ADNI)^{**} cohort. This benefit grew more pronounced after four years, with a reduction of 1.75 points. Similarly, when benchmarked against the expected decline in the BioFINDER cohort, lecanemab showed a reduction of 1.40 points at three years and an even greater reduction of 2.17 points at the four-year mark. In Clarity AD, the most common adverse events ($>10\%$) in the lecanemab group were infusion reactions, ARIA-H (combined cerebral microhemorrhages, cerebral macrohemorrhages, and superficial siderosis), ARIA-E (edema/effusion), headache, and fall.

^{*}The Centiloid scale is used for amyloid PET, where 0 CL is anchored as the average amyloid in young people without amyloid plaques, and 100 is anchored as the average amyloid level in moderate AD. The baseline centiloid level in CLARITY AD was approximately 78 CL. Plaque negativity is defined as conversion to amyloid PET negative (<30 centiloid, or CL).

^{**}Prespecified subgroup analysis of reduced risk of progression: Progression was defined as CDR-SB score progressing to moderate or severe dementia (≥ 9.5), based on Kaplan-Meier plots.

2. About Protofibrils

Protofibrils are thought to be the most toxic A β species that contribute to brain damage in AD and play a major role in the cognitive decline of this progressive and devastating disease. Protofibrils can cause neuronal and synaptic damage in the brain, which can subsequently adversely affect cognitive function through multiple mechanisms.² The mechanism by which this occurs has been reported not only by increasing the formation of insoluble A β plaques, but also by directly damaging signaling between neurons and other cells. It is believed that reducing protofibrils may reduce neuronal damage and cognitive impairment, potentially preventing the progression of AD.³

3. About the Collaboration between Eisai and Biogen for AD

Eisai and Biogen have been collaborating on the joint development and commercialization of AD treatments since 2014. Eisai serves as the lead of lecanemab development and regulatory submissions globally with both companies co-commercializing and co-promoting the product and Eisai having final decision-making authority.

4. About the Collaboration between Eisai and BioArctic for AD

Since 2005, Eisai and BioArctic have had a long-term collaboration regarding the development and commercialization of AD treatments. Eisai obtained the global rights to study, develop, manufacture and market lecanemab for the treatment of AD pursuant to an agreement with BioArctic in December 2007. The development and commercialization agreement on the antibody lecanemab back-up was signed in May 2015.

5. About Eisai Co., Ltd.

Eisai's Corporate Concept is "to give first thought to patients and people in the daily living domain, and to increase the benefits that health care provides." Under this Concept (also known as *human health care (hhc)* Concept), we aim to effectively achieve social good in the form of relieving anxiety over health and reducing health disparities. With a global network of R&D facilities, manufacturing sites and marketing subsidiaries, we strive to create and deliver innovative products to target diseases with high unmet medical needs, with a particular focus in our strategic areas of Neurology and Oncology.

In addition, we demonstrate our commitment to the elimination of neglected tropical diseases (NTDs), which is a target (3.3) of the United Nations Sustainable Development Goals (SDGs), by working on various activities together with global partners.

For more information about Eisai, please visit www.eisai.com (for global headquarters: Eisai Co., Ltd.), and connect with us on [X](#), [LinkedIn](#) and [Facebook](#). The website and social media channels are intended for audiences outside of the UK and Europe. For audiences based in the UK and Europe, please visit www.eisai.eu and Eisai EMEA [LinkedIn](#).

6. About Biogen

Founded in 1978, Biogen is a leading biotechnology company that pioneers innovative science to deliver new medicines to transform patient's lives and to create value for shareholders and our communities. We apply deep understanding of human biology and leverage different modalities to advance first-in-class treatments or therapies that deliver superior outcomes. Our approach is to take bold risks, balanced with return on investment to deliver long-term growth.

The company routinely posts information that may be important to investors on its website at www.biogen.com. Follow Biogen on social media – [Facebook](#), [LinkedIn](#), [X](#), [YouTube](#).

Biogen Safe Harbor

This news release contains forward-looking statements, including about the potential clinical effects of lecanemab; the potential benefits, safety and efficacy of lecanemab; potential regulatory discussions, submissions and approvals and the timing thereof including for lecanemab-irmb (LEQEMBI IQLIK); the treatment of Alzheimer's disease; the anticipated benefits and potential of Biogen's collaboration arrangements with Eisai; the potential of

Biogen's commercial business and pipeline programs, including lecanemab; and risks and uncertainties associated with drug development and commercialization. These forward-looking statements may be accompanied by such words as "aim," "anticipate," "assume," "believe," "contemplate," "continue," "could," "estimate," "expect," "forecast," "goal," "guidance," "hope," "intend," "may," "objective," "plan," "possible," "potential," "predict," "project," "prospect," "should," "target," "will," "would," and other words and terms of similar meaning. Drug development and commercialization involve a high degree of risk, and only a small number of research and development programs result in commercialization of a product. Results in early-stage clinical trials may not be indicative of full results or results from later stage or larger scale clinical trials and do not ensure regulatory approval. You should not place undue reliance on these statements. Given their forward-looking nature, these statements involve substantial risks and uncertainties that may be based on inaccurate assumptions and could cause actual results to differ materially from those reflected in such statements. These forward-looking statements are based on management's current beliefs and assumptions and on information currently available to management. Given their nature, we cannot assure that any outcome expressed in these forward-looking statements will be realized in whole or in part. We caution that these statements are subject to risks and uncertainties, many of which are outside of our control and could cause future events or results to be materially different from those stated or implied in this document, including, among others, uncertainty of long-term success in developing, licensing, or acquiring other product candidates or additional indications for existing products; expectations, plans and prospects relating to product approvals, approvals of additional indications for our existing products, sales, pricing, growth, reimbursement and launch of our marketed and pipeline products; our ability to effectively implement our corporate strategy; the successful execution of our strategic and growth initiatives, including acquisitions; the risk that positive results in a clinical trial may not be replicated in subsequent or confirmatory trials or success in early stage clinical trials may not be predictive of results in later stage or large scale clinical trials or trials in other potential indications; risks associated with clinical trials, including our ability to adequately manage clinical activities, unexpected concerns that may arise from additional data or analysis obtained during clinical trials, regulatory authorities may require additional information or further studies, or may fail to approve or may delay approval of our drug candidates; the occurrence of adverse safety events, restrictions on use with our products, or product liability claims; and any other risks and uncertainties that are described in other reports we have filed with the U.S. Securities and Exchange Commission.

These statements speak only as of the date of this press release and are based on information and estimates available to us at this time. Should known or unknown risks or uncertainties materialize or should underlying assumptions prove inaccurate, actual results could vary materially from past results and those anticipated, estimated or projected. Investors are cautioned not to put undue reliance on forward-looking statements. A further list and description of risks, uncertainties and other matters can be found in our Annual Report on Form 10-K for the fiscal year ended December 31, 2024 and in our subsequent reports on Form 10-Q and Form 10-K, in each case including in the sections thereof captioned "Note Regarding Forward-Looking Statements" and "Item 1A. Risk Factors," and in our subsequent reports on Form 8-K. Except as required by law, we do not undertake any obligation to publicly update any forward-looking statements whether as a result of any new information, future events, changed circumstances or otherwise.

Digital Media Disclosure

From time to time, we have used, or expect in the future to use, our investor relations website (investors.biogen.com), the Biogen LinkedIn account ([linkedin.com/company/biogen-](https://www.linkedin.com/company/biogen/)) and the Biogen X account (<https://x.com/biogen>) as a means of disclosing information to the public in a broad, non-exclusionary manner, including for purposes of the SEC's Regulation Fair Disclosure (Reg FD). Accordingly, investors should monitor our investor relations website and these social media channels in addition to our press releases, SEC filings, public conference calls and websites, as the information posted on them could be material to investors.

References

1. National Institute on Aging (NIA), National Institutes of Health. *What Is Mild Cognitive Impairment?* Available at: <https://www.nia.nih.gov/health/memory-loss-and-forgetfulness/what-mild-cognitive-impairment>. Accessed July 22, 2026.
2. Amin L, Harris DA. A β receptors specifically recognize molecular features displayed by fibril ends and neurotoxic oligomers. *Nat Commun*. 2021;12:3451. doi:10.1038/s41467-021-23507-z.
3. Ono K, Tsuji M. Protofibrils of Amyloid- β are Important Targets of a Disease-Modifying Approach for Alzheimer's Disease. *Int J Mol Sci*. 2020;21(3):952. doi: 10.3390/ijms21030952. PMID: 32023927; PMCID: PMC7037706.